


bsm-52235R**[Primary Antibody]****Bioss**
ANTIBODIES

www.bioss.com.cn


sales@bioss.com.cn

techsupport@bioss.com.cn


400-901-9800

STAT3 Recombinant Rabbit mAb**DATASHEET****Host:** Rabbit**Isotype:** IgG**Clonality:** Recombinant**CloneNo.:** 6G5**GeneID:** 6774**SWISS:** P40763**Target:** STAT3**Immunogen:** A synthesized peptide derived from human STAT3: 1-33.**Purification:** affinity purified by Protein A**Concentration:** 1mg/ml**Storage:** 0.01M TBS (pH7.4) with 1% BSA, 0.02% Proclin300 and 50% Glycerol.
Shipped at 4°C. Store at -20°C for one year. Avoid repeated freeze/thaw cycles.**Background:** The protein encoded by this gene is a member of the STAT protein family. In response to cytokines and growth factors, STAT family members are phosphorylated by the receptor associated kinases, and then form homo- or heterodimers that translocate to the cell nucleus where they act as transcription activators. This protein is activated through phosphorylation in response to various cytokines and growth factors including IFNs, EGF, IL5, IL6, HGF, LIF and BMP2. This protein mediates the expression of a variety of genes in response to cell stimuli, and thus plays a key role in many cellular processes such as cell growth and apoptosis. The small GTPase Rac1 has been shown to bind and regulate the activity of this protein. PIAS3 protein is a specific inhibitor of this protein. Mutations in this gene are associated with infantile-onset multisystem autoimmune disease and hyper-immunoglobulin E syndrome. Alternative splicing results in multiple transcript variants encoding distinct isoforms. [provided by RefSeq, Sep 2015]**Applications:** WB (1:1000-5000)**IHC-P** (1:100-500)**IHC-F** (1:100-500)**IF** (1:100-500)**Flow-Cyt** (1:50-100)**ICC/IF** (1:50-200)**Reactivity:** Human, Mouse, Rat**Predicted MW.:** 88 kDa**Observed MW.:** 85 kDa**Subcellular Location:** Cytoplasm ,Nucleus**VALIDATION IMAGES**


25 ug total protein per lane of various lysates (see on figure) probed with STAT3 monoclonal antibody, unconjugated (bsm-52235R) at 1:1000 dilution and 4°C overnight incubation. Followed by conjugated secondary antibody incubation at r.t. for 60 min.




25 ug total protein per lane of various lysates (see on figure) probed with STAT3 monoclonal antibody, unconjugated (bsm-52235R) at 1:5000 dilution and 4°C overnight incubation. Followed by conjugated secondary antibody incubation at r.t. for 60 min.



Paraformaldehyde-fixed, paraffin embedded Human Kidney; Antigen retrieval by boiling in sodium citrate buffer (pH6.0) for 15 min; Antibody incubation with STAT3 Monoclonal Antibody, Unconjugated (bsm-52235R) at 1:200 overnight at 4°C, followed by conjugation to the bs-0295G-HRP and DAB (C-0010) staining.



Paraformaldehyde-fixed, paraffin embedded Human Kidney Cancer; Antigen retrieval by boiling in sodium citrate buffer (pH6.0) for 15

Blank control: Hela. Primary Antibody (green line): Rabbit Anti-Stat3 antibody (bsm-52235R)
Dilution: 1μg /10⁶ cells; Isotype Control

Important Note: This product as supplied is intended for research use only, not for use in human, therapeutic or diagnostic applications.

min; Antibody incubation with STAT3 Monoclonal Antibody, Unconjugated(bsm-52235R) at 1:200 overnight at 4°C, followed by conjugation to the bs-0295G-HRP and DAB (C-0010) staining.

Antibody (orange line): Rabbit IgG . Secondary Antibody : Goat anti-rabbit IgG-BF488 Dilution: 1µg /test. Protocol The cells were fixed with 4% PFA (10min at room temperature)and then permeabilized with 90% ice-cold methanol for 20 min at -20°C. The cells were then incubated in 5%BSA to block non-specific protein-protein interactions for 30 min at at room temperature .Cells stained with Primary Antibody for 30 min at room temperature. The secondary antibody used for 40 min at room temperature. Acquisition of 20,000 events was performed.

— SELECTED CITATIONS —

- **[IF=8.3]** Qiao Xu. et al. Biological activity study of a pine pollen extract rich in dihydroquercetin for the treatment of psoriasis. PHYTOMEDICINE. 2025 Nov;;157597 WB ;Mouse. 10.1016/j.phymed.2025.157597
- **[IF=8.2]** Qingming Bao. et al. Tumor targeted siRNA delivery by adenosine receptor-specific curdlan nanoparticles. INT J BIOL MACROMOL. 2023 Dec;253:126845 WB ;Mouse. 37703972
- **[IF=5.696]** Jiang-Tao Fan. et al. Exosomal lncRNA NEAT1 from cancer-associated fibroblasts facilitates endometrial cancer progression via miR-26a/b-5p-mediated STAT3/YKL-40 signaling pathway. Neoplasia. 2021 Jul;23:692 WB ;Human. 34153644
- **[IF=5.195]** Yong Wang. et al. An integrated network pharmacology approach reveals that Darutigenol reduces inflammation and cartilage degradation in a mouse collagen-induced arthritis model by inhibiting the JAK-STAT3 pathway. J ETHNOPHARMACOL. 2023 May;;116574 WB ;Mouse. 37160212
- **[IF=4.7]** Ya-Zhao Zhang. et al. Exploring the alleviating effect and mechanism of the diterpenoid alkaloid Polyschistine D on rheumatoid arthritis based on network pharmacology and non-targeted metabolomics. INT IMMUNOPHARMACOL. 2025 Nov;165:115436 WB ;Mouse. 40907334