bs-1313R **[Primary Antibody]**

VEGFA | VEGF Rabbit pAb

www.bioss.com.cn
sales@bioss.com.cn
techsupport@bioss.com.cn
400-901-9800

DATASHEET

Host: Rabbit **Isotype:** IgG

Clonality: Polyclonal

GeneID: 7422 **SWISS:** P15692

Target: VEGFA | VEGF

Immunogen: KLH conjugated synthetic peptide derived from human VEGF: 102-213/213.

Purification: affinity purified by Protein A

Concentration: 1mg/ml

Storage: 0.01M TBS (pH7.4) with 1% BSA, 0.02% Proclin300 and 50% Glycerol.
Shipped at 4°C. Store at -20°C for one year. Avoid repeated freeze/thaw cycles.

Background: Vascular endothelial growth factor (VEGF), originally known as vascular permeability factor (VPF), is a signal protein produced by cells that stimulates the formation of blood vessels. To be specific, VEGF is a sub-family of growth factors, the platelet-derived growth factor family of cystine-knot growth factors. They are important signaling proteins involved in both vasculogenesis (the de novo formation of the embryonic circulatory system) and angiogenesis (the growth of blood vessels from pre-existing vasculature).

Applications: **WB** (1:500-2000)
IHC-P (1:100-500)
IHC-F (1:100-500)
IF (1:100-500)


Reactivity: Human, Mouse, Rat, Rabbit (predicted: Pig, Cow, Chicken, Dog)

Predicted MW.: 23 kDa


Observed MW.: 27 kDa

Subcellular Location: Secreted


VALIDATION IMAGES




Sample:Kidney(Mouse) Lysate at 30 ug Primary: Anti-VEGF (bs-1313R) at 1/300 dilution Secondary: IRDye800CW Goat Anti-Rabbit IgG at 1/20000 dilution Predicted band size: 23 kD Observed band size: 27 kD




Tissue/cell: rabbit liver tissue; 4% Paraformaldehyde-fixed and paraffin-embedded; Antigen retrieval: citrate buffer (0.01M, pH 6.0), Boiling bathing for 15min; Block endogenous peroxidase by 3% Hydrogen peroxide for 30min; Blocking buffer (normal goat serum,C-0005) at 37∩ for 20 min; Incubation: Anti-VEGF Polyclonal Antibody, Unconjugated(bs-1313R) 1:400, overnight at 4ΣC, followed by conjugation to the secondary antibody(SP-0023) and DAB(C-0010) staining



Tissue/cell: rat brain tissue; 4% Paraformaldehyde-fixed and paraffin-embedded; Antigen retrieval: citrate buffer (0.01M, pH 6.0), Boiling bathing for 15min; Block endogenous peroxidase by 3% Hydrogen peroxide for 30min; Blocking buffer (normal goat serum,C-0005) at 37∩ for 20 min; Incubation: Anti-VEGF Polyclonal Antibody, Unconjugated(bs-1313R) 1:300, overnight at 4ΣC, followed by conjugation to the secondary antibody(SP-0023) and DAB(C-0010) staining



Paraformaldehyde-fixed, paraffin embedded (rat kidney); Antigen retrieval by boiling in sodium citrate buffer (pH6.0) for 15min; Blocking buffer (normal goat serum) at 37°C for 30min; Incubation with (VEGFA) Polyclonal Antibody, Unconjugated (bs-1313R) at 1:200 overnight at 4°C, followed by a conjugated Goat Anti-Rabbit



Paraformaldehyde-fixed, paraffin embedded (mouse kidney); Antigen retrieval by boiling in sodium citrate buffer (pH6.0) for 15min; Blocking buffer (normal goat serum) at 37°C for 30min; Incubation with (VEGFA) Polyclonal Antibody, Unconjugated (bs-1313R) at 1:200 overnight at 4°C, followed by a conjugated Goat Anti-Rabbit

Important Note: This product as supplied is intended for research use only, not for use in human, therapeutic or diagnostic applications.

IgG antibody (bs-0295G-BF594) for 90 minutes, and DAPI for nuclei staining.

IgG antibody (bs-0295G-BF594) for 90 minutes, and DAPI for nuclei staining.

SELECTED CITATIONS

- **[IF=21.3]** Liu Haoran. et al. A Bioflexible Janus Dressing with Unidirectional Drainage for Dynamic pH Monitoring, Exudate Management, and Immune Modulation in Diabetic Wound Healing. ADV FIBER MATER. 2026 Mar;:1-15 IHC ;Rat. 10.1007/s42765-025-00661-3
- **[IF=19]** Yaning Qu. et al. Microenvironment-Regulated Dual-Layer Microneedle Patch for Promoting Periodontal Soft and Hard Tissue Regeneration in Diabetic Periodontitis. ADV FUNCT MATER. 2025 Jan;:2418076 IHC ;Rat. 10.1002/adfm.202418076
- **[IF=18.027]** Guanghao Wu. et al. Enhanced Proliferation of Visualizable Mesenchymal Stem Cell–Platelet Hybrid Cell for Versatile Intracerebral Hemorrhage Treatment. ACS NANO. 2023;XXXX(XXX):XXX-XXX IF ;Mouse. 37037487
- **[IF=18]** Shan He. et al. A nanoenzyme-modified hydrogel targets macrophage reprogramming-angiogenesis crosstalk to boost diabetic wound repair. BIOACT MATER. 2024 May;35:17 IF ;Mouse. 38304915
- **[IF=17.521]** Huan Lei. et al. A Combination Therapy Using Electrical Stimulation and Adaptive, Conductive Hydrogels Loaded with Self-Assembled Nanogels Incorporating Short Interfering RNA Promotes the Repair of Diabetic Chronic Wounds. Advanced Science. 2022 Sep;:2201425 IF ;Rat. 36064844

Important Note: This product as supplied is intended for research use only, not for use in human, therapeutic or diagnostic applications.