bs-1063R **[Primary Antibody]**

ADRB3 Rabbit pAb

www.bioss.com.cn
sales@bioss.com.cn
techsupport@bioss.com.cn
400-901-9800

DATASHEET

Host: Rabbit **Isotype:** IgG

Clonality: Polyclonal

GeneID: 155 **SWISS:** P13945

Target: ADRB3

Immunogen: KLH conjugated synthetic peptide derived from human ADRB3: 310-408/408. < Cytoplasmic >

Purification: affinity purified by Protein A

Concentration: 1mg/ml

Storage: 0.01M TBS (pH7.4) with 1% BSA, 0.02% Proclin300 and 50% Glycerol.
Shipped at 4°C. Store at -20°C for one year. Avoid repeated freeze/thaw cycles.

Background: beta 3 Adrenergic Receptor is located mainly in adipose tissue and is involved in the regulation of lipolysis and thermogenesis. Beta adrenergic receptors are involved in the epenephrine and norepinephrine-induced activation of adenylate cyclase through the action of G proteins. beta 3 Adrenergic Receptor expression has been documented in adipose tissues, heart, and in smooth muscle of digestive and urinary tract organs (bladder, colon, small intestine, stomach, ureter). Utilization of alternate promoters and/or 3-prime untranslated regions may result in tissue-specific regulation of the expression of this protein.


Applications: **WB** (1:500-2000)
IHC-P (1:100-500)
IHC-F (1:100-500)
IF (1:100-500)

Reactivity: Human, Rat (predicted: Mouse, Pig, Sheep, Dog)


Predicted MW.: 44 kDa

Subcellular Location: Cell membrane

VALIDATION IMAGES



Sample: K562(Human) Cell Lysate at 40 ug Primary: Anti-beta 3 Adrenergic Receptor (bs-1063R) at 1/300 dilution Secondary: IRDye800CW Goat Anti-Rabbit IgG at 1/20000 dilution Predicted band size: 44 kD Observed band size: 54 kD



Tissue/cell: Rat adrenal gland; 4% Paraformaldehyde-fixed and paraffin-embedded; Antigen retrieval: citrate buffer (0.01M, pH 6.0), Boiling bathing for 15min; Block endogenous peroxidase by 3% Hydrogen peroxide for 30min; Blocking buffer (normal goat serum,C-0005) at 37°C for 20 min; Incubation: Anti-ADRB3 Polyclonal Antibody, Unconjugated(bs-1063R) 1:500, overnight at 4°C, followed by conjugation to the secondary antibody(SP-0023) and DAB(C-0010) staining

SELECTED CITATIONS

- **[IF=15.7]** Sánchez-Feutrie Manuela. et al. Identification of Neuritin 1 as a local metabolic regulator of brown adipose tissue. NAT COMMUN. 2025 Sep;16(1):1-18 WB ;Mouse. 40908286
- **[IF=6.208]** Lina S. Farhoumand. et al. Blockade of ß-Adrenergic Receptors by Nebivolol Enables Tumor Control Potential for Uveal Melanoma in 3D Tumor Spheroids and 2D Cultures. INT J MOL SCI. 2023 Jan;24(6):5894 FCM ;Human. 36982966
- **[IF=4.757]** Katarina Andelova. et al. Cardiac Cx43 Signaling Is Enhanced and TGF-β1/SMAD2/3 Suppressed in Response

Important Note: This product as supplied is intended for research use only, not for use in human, therapeutic or diagnostic applications.

to Cold Acclimation and Modulated by Thyroid Status in Hairless SHRM. BIOMEDICINES. 2022 Jul;10(7):1707 WB ;Rat. 35885012

- **[IF=4.699]** Pellegrino Lippiello. et al. Role of β3‐adrenergic receptor in the modulation of synaptic transmission and plasticity in mouse cerebellar cortex. J Neurosci Res. 2020 Nov;98(11):2263-2274 IF ;Mouse. 33174240
- **[IF=4.2]** Wang Yu-Wan. et al. Long-term exposure to constant light disrupts intestinal stem cells through sympathoexcitation-induced Wnt5a signaling inhibition. GASTROENTEROL REP. 2025 May;13: IHC ;Rat. 40438259

Important Note: This product as supplied is intended for research use only, not for use in human, therapeutic or diagnostic applications.