

bsm-60866R**[Primary Antibody]****BioSS**
ANTIBODIES

www.bioss.com.cn


sales@bioss.com.cn

techsupport@bioss.com.cn

400-901-9800

Citrulline-Histone H3 (Arg2/8/17) Recombinant Rabbit mAb**— DATASHEET —**

Host: Rabbit	Isotype: IgG	Applications: WB (1:2000-20000) IHC-P (1:200-800) IHC-F (1:200-800) IF (1:200-800) Reactivity: Human (predicted: Mouse, Rat) Predicted MW.: 15 kDa Subcellular Location: Nucleus
Clonality: Recombinant	CloneNo.: 11E7	
Target: Citrulline-Histone H3 (Arg2/8/17)		
Purification: affinity purified by Protein A		
Concentration: 1mg/ml		
Storage: PBS, Glycerol, BSA. Shipped at 4°C. Store at -20°C for one year. Avoid repeated freeze/thaw cycles.		
Background: Modulation of the chromatin structure plays an important role in the regulation of transcription in eukaryotes. The nucleosome, made up of four core histone proteins (H2A, H2B, H3 and H4), is the primary building block of chromatin. The N-terminal tail of core histones undergoes different posttranslational modifications including acetylation, phosphorylation and methylation. These modifications occur in response to cell signal stimuli and have a direct effect on gene expression. In most species, the histone H2B is primarily acetylated at lysines 5, 12, 15 and 20. Histone H3 is primarily acetylated at lysines 9, 14, 18 and 23. Acetylation at lysine 9 appears to have a dominant role in histone deposition and chromatin assembly in some organisms. Phosphorylation at Ser10 of histone H3 is tightly correlated with chromosome condensation during both mitosis and meiosis.		

— VALIDATION IMAGES —

Paraformaldehyde-fixed, paraffin embedded Human Ovarian Cancer; Antigen retrieval by boiling in sodium citrate buffer (pH6.0) for 15 min; The section was incubated with Citrulline-Histone H3 (Arg2/8/17) Monoclonal Antibody, Unconjugated (bsm-60866R) at 1:200 overnight at 4°C, followed by conjugation to the bs-0295G-HRP and DAB (C-0010) staining.

— SELECTED CITATIONS —

- **[IF=5]** Haoyun Luo. et al. STING Drives Psoriatic Inflammation by Promoting Neutrophil Recruitment and Facilitating NETosis. IMMUNOLOGY. 2026 Mar;; WB ;Mouse. 41775627