
Smad1 Recombinant Rabbit mAb

Catalog Number: bsm-52222R

Target Protein: Smad1

Concentration: 1mg/ml

Form: Liquid

Host: Rabbit

Clonality: Recombinant

Clone No.: 2D9

Isotype: IgG

Applications: WB (1:1000-2000), IHC-P (1:100-500), IHC-F (1:50-200), ICC/IF (1:50-200), IF (1:50-200)

Reactivity: Human (predicted:Mouse)

Predicted MW: 52 kDa

Entrez Gene: 4086

Swiss Prot: Q15797


Purification: affinity purified by Protein A

Storage: 0.01M TBS (pH7.4) with 1% BSA, 0.02% Proclin300 and 50% Glycerol.

Shipped at 4°C. Store at -20°C for one year. Avoid repeated freeze/thaw cycles.

Background: The protein encoded by this gene belongs to the SMAD, a family of proteins similar to the gene products of the Drosophila gene 'mothers against decapentaplegic' (Mad) and the C. elegans gene Sma. SMAD proteins are signal transducers and transcriptional modulators that mediate multiple signaling pathways. This protein mediates the signals of the bone morphogenetic proteins (BMPs), which are involved in a range of biological activities including cell growth, apoptosis, morphogenesis, development and immune responses. In response to BMP ligands, this protein can be phosphorylated and activated by the BMP receptor kinase. The phosphorylated form of this protein forms a complex with SMAD4, which is important for its function in the transcription regulation. This protein is a target for SMAD-specific E3 ubiquitin ligases, such as SMURF1 and SMURF2, and undergoes ubiquitination and proteasome-mediated degradation. Alternatively spliced transcript variants encoding the same protein have been observed. [provided by RefSeq].

VALIDATION IMAGES



Sample: Lane 1: Human Huvec cell lysates Lane 2: Human HeLa cell lysates Lane 3: Human MCF-7 cell lysates
 Primary: Anti-Smad1 (bsm-52222R) at 1/1000 dilution Secondary: IRDye800CW Goat Anti-Rabbit IgG at 1/20000 dilution Predicted band size: 52 kDa Observed band size: 54 kDa