bs-1461R **[Primary Antibody]**

VDAC1 Rabbit pAb, Loading Control

www.bioss.com.cn
sales@bioss.com.cn
techsupport@bioss.com.cn
400-901-9800

DATASHEET

Host: Rabbit **Isotype:** IgG

Clonality: Polyclonal

GeneID: 7416

Target: VDAC1

Immunogen: KLH conjugated synthetic peptide derived from human VDAC: 85-190/283.

Purification: affinity purified by Protein A

Concentration: 1mg/ml

Storage: 0.01M TBS (pH7.4) with 1% BSA, 0.02% Proclin300 and 50% Glycerol.
Shipped at 4°C. Store at -20°C for one year. Avoid repeated freeze/thaw cycles.

Background: Voltage dependent anion selective channel protein 1 (VDAC/Porin) belongs to the eukaryotic mitochondrial porin family and forms a channel through the mitochondrial outer membrane and also the plasma membrane. The channel allows diffusion of small hydrophilic molecules; it adopts an open conformation at low or zero membrane potential and a closed conformation at potentials above 30-40 mV. The open state has a weak anion selectivity whereas the closed state is cation selective. VDAC/Porin expression is observed in the heart,liver and skeletal muscle.

Applications: **WB** (1:500-2000)
IHC-P (1:100-500)
IHC-F (1:100-500)
IF (1:100-500)


Reactivity: Human, Rat (predicted: Mouse, Rabbit, Pig, Sheep, Cow, Dog, Horse)

Predicted MW.: 32 kDa


Observed MW.: 25 kDa

Subcellular Location: Cell membrane ,Cytoplasm

VALIDATION IMAGES



Sample: Jurkat(Human) Cell Lysate at 30 ug
Primary: Anti-VDAC (bs-1461R) at 1/500 dilution
Secondary: IRDye800CW Goat Anti-Rabbit IgG at 1/20000 dilution Predicted band size: 32 kD
Observed band size: 25 kD



Tissue/cell: rat brain tissue; 4% Paraformaldehyde-fixed and paraffin-embedded; Antigen retrieval: citrate buffer (0.01M, pH 6.0), Boiling bathing for 15min; Block endogenous peroxidase by 3% Hydrogen peroxide for 30min; Blocking buffer (normal goat serum,C-0005) at 37°C for 20 min; Incubation: Anti-VDAC Polyclonal Antibody, Unconjugated(bs-1461R) 1:200, overnight at 4°C, followed by conjugation to the secondary antibody(SP-0023) and DAB(C-0010) staining

SELECTED CITATIONS

- **[IF=7.69]** Manczak Maria. et al. Abnormal interaction of VDAC1 with amyloid beta and phosphorylated tau causes mitochondrial dysfunction in Alzheimer's disease. HUM MOL GENET. 2012 Aug;21(23):5131-5146 WB,IP ;="Human, Mouse". 22926141
- **[IF=6.575]** Qiyu Gan. et al. Prognostic Value and Immune Infiltration of HPV-Related Genes in the Immune Microenvironment of Cervical Squamous Cell Carcinoma and Endocervical Adenocarcinoma. CANCERS. 2023 Jan;15(5):1419 IHC ;Human. 36900213
- **[IF=5.19]** Peng Junjun. et al. Mitochondria-associated endoplasmic reticulum membranes participate mitochondrial

Important Note: This product as supplied is intended for research use only, not for use in human, therapeutic or diagnostic applications.

dysfunction and endoplasmic reticulum stress caused by copper in duck kidney. ENVIRON SCI POLLUT R. 2023 May;:1-12 WB ;Duck. 37253910

- **[IF=4.1]** Yuan Li. et al. Pestivirus bovine viral diarrhea virus infection induces ROS–HIF-1a axis-driven glycolytic reprogramming, which increases viral replication by impairing RIG-I-dependent type I interferon response. J VIROL. 2026 May WB ;Cow. 42095650
- **[IF=2.73]** Wang, Zhaoqi, et al. "Penehyclidine hydrochloride prevents anoxia/reoxygenation injury and induces H9c2 cardiomyocyte apoptosis via a mitochondrial pathway." European Journal of Pharmacology (2017). WB ;="Rat". 28089921

Important Note: This product as supplied is intended for research use only, not for use in human, therapeutic or diagnostic applications.