

bs-2226G**[Secondary Antibodies]****Bioss**
ANTIBODIES

www.bioss.com.cn

sales@bioss.com.cn

techsupport@bioss.com.cn


400-901-9800

Goat anti-Human IgE pAb**— DATASHEET —****Host:** Goat**Isotype:** IgG**Clonality:** Polyclonal**Target:** Goat anti-Human IgE pAb**Purification:** affinity purified by Protein G**Storage:** 0.01M PBS (pH7.4).

Shipped at 4°C. Store at -20°C. Avoid repeated freeze/thaw cycles.

Background: IgE is the least abundant immunoglobulin in plasma, found at a concentration of less than 0.6 micrograms/ml of normal plasma. Elevated IgE levels are found in patients experiencing severe allergic reactions and parasitic infections. In a myeloma condition, IgE is produced by a single clone of plasma cells. The structure of myeloma IgE, however, is normal, and the immunoglobulin purified from a myeloma source is a useful protein for studying immunoglobulin behavior. The affinity purified IgE reacted only with anti IgE and not with anti IgG, IgA, IgM or IgD by immunodiffusion and IEP techniques. Prepared from myeloma plasma shown to be non reactive for HbsAG, anti-HCV, anti-HBc, and negative for anti-HIV 1 & 2 by FDA approved tests.

Applications: Isotype Control
Blocking Assay
etc.
Conjugate-Dependent.

Reactivity: Human**— VALIDATION IMAGES —**

100 ng Human IgE protein (bs-2226P) per lane probed with Human IgE polyclonal antibody respectively, unconjugated (bs-2226G) at 1:1000 dilution and 4°C overnight incubation. Followed by corresponding conjugated secondary antibody incubation at r.t. for 60 min.