

bs-0375R

[Secondary Antibodies]



www.bioss.com.cn

sales@bioss.com.cn

techsupport@bioss.com.cn

400-901-9800

Rabbit Anti-Guinea Pig IgM

— DATASHEET —

Host: Rabbit Clonality: Polyclonal Target: Rabbit Anti-Guinea Pig IgM Immunogen: Native Guinea pig IgM: full length. Purification: affinity purified by Protein A Storage: 0.01M PBS (pH7.4). Store at -20°C stable for 2 years (lyophilized powder). Avoid repeated freeze/thaw cycles. Background: Immunoglobulin M (IgM) normally constitutes about 10% of serum immunoglobulins. IgM antibody is prominent in early immune responses to most antigens and is largely confined to plasma due to its large size. Monomeric IgM is expressed as a membrane bound antibody on the surface of B cells and as a pentamer when secreted by plasma cells. Due to its high valency IgM is more efficient than other isotypes in binding antigens with repeating epitopes (virus particles and red blood cells) and is more efficient than IgG in activating the complement pathway. The gene for the mu constant region contains four domains separated by short intervening sequences.	Isotype: IgG Applications: Isotype Control Blocking Assay etc. Conjugate-Dependent. Reactivity: Guinea Pig
--	--

— SELECTED CITATIONS —

- **[IF=4.5]** Yingjun Cui. et al. Salp14 epitope-based mRNA vaccination induces early recognition of a tick bite. VACCINE. 2024 Oct;42:126304 ELISA ;Guinea pig. 39236403